



Nuts and diabetes management: Nutritional and metabolic benefits

Nomagugu Ndlovu

Department of Biological and Environmental Science, Faculty of Natural Sciences, Walter Sisulu University, Nelson Mandela Drive, P Bag X1, Mthatha, South Africa

Corresponding author: Nomagugu Ndlovu, PhD, Department of Biological and Environmental Science, Faculty of Natural Sciences, Walter Sisulu University, Nelson Mandela Drive, P Bag X1, Mthatha, South Africa

Submission Date: May 6th, 2025; **Acceptance Date:** August 21st, 2025; **Publication Date:** August 22nd, 2025

Please cite this article as: Ndlovu N. Nuts and diabetes management: Nutritional and metabolic benefits.

Bioactive Compounds in Health and Disease 2025; 8(8): 243 – 258. DOI: <https://doi.org/10.31989/bchd.8i8.1635>

ABSTRACT


The escalating global prevalence of diabetes has spurred the urgency for effective management strategies. In recent years, there has been growing interest in the potential role of dietary interventions, particularly the incorporation of nuts, in the management of diabetes.

This review aims to elucidate the nutritional significance of nuts and their potential impact on diabetes management. Articles used in this narrative review were retrieved from Scopus, Google scholar, ResearchGate, and other research platforms using keywords/phrases such as 'diabetes', 'diabetes management' 'dietary strategies for diabetes' as well as 'nuts in diabetes management'. Only articles written in English, peer-reviewed, and focused on dietary strategies for managing diabetes were included in this narrative review.

Findings showed that nuts, abundant in unsaturated fats, fiber, protein, vitamins, and minerals, exhibit favourable effects on glycaemic regulation, insulin sensitivity and lipid profiles. Accumulated evidence from epidemiological studies, clinical trials, and meta-analyses suggests that habitual nut consumption holds promise in ameliorating glycaemic control, mitigating cardiovascular complications, and augmenting overall health outcomes among individuals with diabetes. However, the specific mechanisms underlying these effects remain to be fully elucidated, and further research is warranted to establish optimal recommendations for nut consumption in the context of diabetes management.

Although incorporating nuts into a balanced diet represents a promising and practical approach for enhancing the nutritional quality and metabolic health of individuals with diabetes, it is essential to address potential allergenicity and safety concerns associated with their consumption.

Keywords: nuts, diabetes, glycaemia, insulin resistance, lipid profile



Graphical Abstract: Nuts and Diabetes Management: Nutritional and Metabolic benefits

©FFC 2025. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0>)

INTRODUCTION

Diabetes mellitus (DM) is a chronic global epidemic, a major health problem, and is one of the fastest-growing global health emergencies of the 21st century [1,2]. Diabetes has emerged as the ninth leading cause of mortality globally [3,4]. Approximately 10.5 million adults were diagnosed with diabetes, with over 90% suffering from Type 2 Diabetes (T2D) [1], making T2D the most common type of diabetes. In 2021, the International Federation of Diabetes reported that 537 million adults are now living with diabetes worldwide, an increase

of 16% (74 million) since the previous IDF estimates in 2019 [4,5]. Projections suggest that this figure will escalate to 783 million or 1 in 8 people by 2045 [4,5]. This would be an increase of 46%, more than double the estimated population growth (20%) over the same period [4]. The rise in the number of people with type 2 diabetes is of socio-economic, demographic, environmental and genetic factors [5]. The key contributors include urbanisation, an ageing population, decreasing levels of physical activity and increasing prevalence of overweight and obesity [5,6].

Diabetes not only elevates blood glucose levels but also triggers numerous long-term complications that significantly impact the health and well-being of individuals affected by it [7]. Of particular concern is its propensity to precipitate cardiovascular disease (CVD), encompassing conditions like coronary artery disease, heart failure, and stroke, which stand as leading causes of mortality and disability among those with diabetes [4,5]. Research consistently demonstrates that individuals with diabetes face a heightened risk, two to four times greater, of developing CVD compared to those without the condition [8,9]. This increased risk stems from various factors such as insulin resistance, dyslipidemia, hypertension, and systemic inflammation, all of which exacerbate atherosclerosis, the primary driver of most cardiovascular incidents [8,10]. Moreover, diabetes exacerbates endothelial dysfunction, oxidative stress, and abnormal platelet function, further amplifying the likelihood of cardiovascular complications [10,11]. Concurrently experiencing both diabetes and CVD compounds the situation, heightening the probability of recurrent episodes and driving up healthcare expenditures [12]. Inadequately managed diabetes can lead to numerous other issues, including kidney problems, nerve damage, eye issues, digestive disorders, erectile dysfunction, skin conditions, increased vulnerability to infections, and dental issues [8,13]. Considering this, it is important to find effective ways to manage the cardiovascular risk in diabetic patients. Lifestyle changes, such as those pertaining to diet, form the foundation of a broad strategy for reducing and managing the risk of diabetes [4,14].

The goal of diabetes treatment is to normalize fasting and postprandial blood glucose and lipids, to attain normal blood pressure, and to maintain a healthy weight [15,16]. Therefore, current guidelines for diabetes management advocate for dietary plans like the Mediterranean and vegetarian diets, which promote the consumption of nuts [17,18]. Nuts, including tree nuts

(like almonds, Brazil nuts, cashews, hazelnuts, macadamias, pecans, pine nuts, pistachios, and walnuts) and peanuts are rich in nutrients and offer complex compositions containing unsaturated fatty acids, plant-based proteins, minerals without sodium, as well as phenolic and other bioactive compounds [14]. This review differs from previous studies by integrating evidence to explain how specific components in nuts improve diabetes and metabolic syndrome, including conditions like poor glycemic control, dyslipidemia, inflammation, and insulin resistance.. The novelty of this paper lies in its synthesis of multi-nutrient mechanisms, from unsaturated fats and fiber to phenolic compounds, and its emphasis on nut-specific bioactivity. By detailing molecular mechanisms such as AMPK activation, GLP-1 secretion, suppression of lipogenic genes, and modulation of gut microbiota, this review highlights how nuts can serve as a functional dietary therapy beyond traditional glycaemic management, positioning them as a holistic intervention for metabolic syndrome.

METHODS

This narrative review assesses the benefits of nut consumption for diabetes and metabolic syndrome.. The literature was sourced from multiple academic databases including Scopus, PubMed, Google Scholar, and ResearchGate, using the following keywords: “nuts,” “diabetes,” “metabolic syndrome,” “glycaemic control,” “insulin resistance,” “lipid profile,” “dietary intervention.” Boolean operators (AND, OR) were applied to refine the search. Only peer-reviewed articles published in English between January 2000 and April 2024 were considered. Inclusion criteria encompassed original research articles, clinical trials, systematic reviews, and meta-analyses that investigated the relationship between nut consumption and glycaemic, lipid, or metabolic outcomes. Studies without nutritional data or mechanistic relevance were excluded. Screening of titles and abstracts was conducted manually by the author. A total of 23 studies

were included in the final synthesis. Key data extracted included study population, nut type and dosage, duration of intervention, and main metabolic outcomes.

Nutrient composition of nuts: Nuts, typically characterized as dry fruits with an edible seed encased in a hard shell, such as cashews, walnuts, almonds, chestnuts, pistachios, and hazelnuts, are globally recognized for their nutritional richness [19,20]. They provide essential nutrients like monounsaturated and polyunsaturated fatty acids, vitamins E and K, as well as minerals such as magnesium, copper, potassium, and selenium, alongside dietary fibers, carotenoids, and phytosterols, which contribute to potential antioxidant benefits [20]. Their compact size also makes them convenient for consumption in various settings. Furthermore, incorporating nuts into the diet is often associated with mitigating risk factors for chronic diseases such as oxidative stress, inflammation, visceral adiposity, hyperglycemia, insulin resistance, endothelial dysfunction, and metabolic syndrome, thanks to their diverse nutrient profile [21,22]. Notably, many of the antioxidants in nuts are concentrated in their outer layers, such as the pellicle in almonds and peanuts [21] implying that when the skin is removed from the nuts the antioxidants may be lost.

Proteins: Nuts are a rich source of proteins and essential amino acids [21,22]. Pistachio has the highest protein content ranging from 19.4-22.1% [22], while chestnut has the lowest protein content (1.63%) [21]. In terms of the amino acid profile found in each type of nut, there is significant variability in both essential and non-essential amino acid content [21]. Variations may occur due to factors like variety and location. Glutamic acid stands out as the most abundant non-essential amino acid, ranging from 0.02 g/100 g in chestnuts to 6.21 g/100 g in almonds

[23]. Arginine follows as the second most abundant non-essential amino acid, ranging from 0.12 g/100 g in chestnuts to 3.08 g/100 g in peanuts, with aspartic acid ranging between 0.03 g/100 g in chestnuts and 3.15 g/100 g in peanuts [21,23]. Among the essential amino acids, leucine is the most prominent, followed by phenylalanine and valine. Chestnuts exhibit the lowest levels of these essential amino acids (0.10 g/100 g, 0.07 g/100 g, and 0.09 g/100 g for leucine, phenylalanine, and valine, respectively), while peanuts are rich in leucine and phenylalanine, and pistachios are particularly high in valine [24]. Pecan can contain up to 17.84% protein content and all the amino acids essential for human diet (isoleucine, leucine, valine, lysine, threonine, tryptophan, phenylalanine, methionine and histidine) [25]. Although nut proteins when compared to animal proteins are often considered incomplete, their consumption is linked to cardiovascular health [26]. Furthermore, the significant presence of arginine in all tree nuts offers benefits for immune response, inflammation, cardiovascular function, and reproductive performance, thereby reducing the risk of cardiovascular disease [21].

Vitamins: Nuts contain fat-soluble vitamins and antioxidants like α -tocopherol (known as vitamin E), which contribute to improved health, combat aging, enhance brain function, and promote healthy skin [21]. Various studies have reported an association between a increased dietary intake of fat-soluble vitamins K and E, with lowered risk of type 2 diabetes [27–29]. Specific nuts like walnuts, almonds, pine nuts, and hazelnuts are notably rich in vitamin E, while almonds, cashews, pistachios, walnuts, and peanuts are abundant in B vitamins [30]. Vitamin B₆ has been shown to protect endothelial function and improve insulin resistance and low Vitamin B₆ status might be a risk factor for non-alcoholic fatty liver disease (NAFLD) [31]. Pistachios and

chestnuts have higher concentrations of folic acid, with chestnuts being the richest in vitamin C [32]. Supplemental vitamin C has been shown to potentially improve glycemic control and BP in people with type 2 diabetes [33,34].

Minerals: Nuts serve as valuable reservoirs of essential minerals like magnesium and potassium. In recent times, the promotion of increased nut consumption has been recognized as beneficial for human health due to their role in augmenting mineral intake [35]. When consumed moderately, nuts are deemed as heart-healthy snacks [35]. They are notably rich in minerals like copper and magnesium, which are believed to offer protective effects against coronary heart disease [21]. Moreover, certain varieties are particularly abundant in potassium, for example, pistachios (642-1025mg/100g) and cashews (660mg/100g)[21]. While most nuts contain notable amounts of zinc and iron, pine nuts (Zn: 3.08-6.45mg/100g; Fe: 5.53-6.64mg/100g), cashews (Zn: 0.96-5.78mg/100g; Fe: 3.82-6.68mg/100g), and almonds (1.91-3.12mg/100g; Fe: 3.71-6.21mg/100g) excel in this regard[21]. Despite not being notably rich in calcium overall, select nuts like almonds exhibit relatively higher calcium content (248mg/100g) [36].

Fiber: Dietary fiber, a vital component of human nutrition, can be categorized into two main types based on their behaviour in aqueous solutions: insoluble and soluble fiber. Insoluble fiber, comprising cellulose, lignin, and various hemicelluloses primarily found in grains and vegetables, serves as a bulking agent in the digestive system, promoting efficient intestinal transit[37]. In contrast, water-soluble dietary fibers encompass a diverse range of polysaccharides such as pectins, gums, mucilages, algal polysaccharides, and certain hemicelluloses and storage polysaccharides[38]. These

fibers exhibit a notable capacity to absorb and retain water, often forming highly viscous solutions upon dissolution[38]. Scientific studies have shown that higher intake of dietary fiber is linked to lower risks of obesity, T2DM, cancer, and cardiovascular disease [39]. Almonds, among all nuts, contain the highest fiber content, influenced by genetic factors. Research indicates that the fiber content in almonds varies, ranging from 6.88% to 9.74%, with almond skin contributing significantly, comprising around 60% of the fiber. Cashews have the lowest fiber content, typically around 3% to 4%, while chestnuts are considered a good fiber source and that of hazelnuts ranges from 6.5 to 9.7%[40]. It is believed that an increase in dietary fiber intake results in a lower pH level in the gut, which increases the solubility of calcium ions there by increasing its absorption [41,42]. Calcium has a vital role in the prevention of diabetes by improving insulin sensitivity and pancreatic β -cell functions[43]. Additionally, fiber has been reported to reduce post prandial glycaemia[41]. The reduction in glucose levels due to fiber occurs by the soluble dietary fiber forming a gel, causing the contents of the gut to become thicker in both the stomach and intestines. This increased thickness slows down the diffusion of nutrients, gastric emptying, and absorption of nutrients in the small intestine[38]. Consequently, carbohydrates are absorbed into the bloodstream at a slower pace, resulting in more stable post prandial glycaemia[44].

Lipids: Dietary fats are divided into three subgroups: unsaturated FA, saturated fatty acids (SFA), and trans fatty acids (TFA)[45]. Unsaturated FA consists of monounsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA) [46]. Although the fatty acid composition of the diet plays a significant role in enhancing insulin sensitivity and reducing T2DM and T2DM-related complications, the underlying mechanisms

remain unclear[45]. It is claimed that saturated and trans fats are possible risk factors of T2DM while the poly-unsaturated fatty acids (PUFAs) may have positive effects on people with T2DM[45]. Nuts serve as an excellent source of fats following vegetable oil seeds [21]. The lipid content in nuts typically ranges from 40% to 75%[39]. Varieties such as walnut, macadamia, pine nut, and pecans tend to have higher lipid content (around 70%) compared to cashew, almond, pistachio, and hazelnut, which generally contain between 45% and 62% lipid[14]. The lipid profiles of different nuts are shown in Table 1. The PUFAs content ranges from 1.5-47.2g/100g, with the walnuts having the highest PUFAs content and

macademia nuts having the least [40]. A diet high in unsaturated fatty acids, in particular, long-chain omega-6 and omega-3 polyunsaturated fatty acids, has been shown to lead to the suppression of lipogenic genes (genes of lipid synthesis) and induction of genes involved with fatty acid oxidation, which may reduce hepatic insulin resistance[14]. Additionally, nuts contain phytosterols which is a component of the cell membrane and is reported to interfere with cholesterol absorption and thus helping lower blood cholesterol when present in sufficient amounts in the intestinal lumen[14,39]. Fatty acid profiles of different nuts are shown in Table 1 below.

Table 1. Fatty acid profiles of different types of nuts.

Nuts	Total Fat	SFA	MUFAs	PUFAs	Omega-6	Omega-3
Almonds	50.6	3.9	32.2	12.2	12.2	0.00
Walnuts	65.2	6.1	8.9	47.2	38.1	9.08
Pistachio	44.4	5.4	23.3	13.5	13.2	0.25
Cashew	46.4	9.2	27.3	7.8	7.7	0.15
Pine nut	68.4	4.9	18.8	34.1	33.2	0.16
Pecan	72.0	6.2	40.8	21.6	20.6	1.00
Macadamia	75.8	12.1	58.9	1.5	1.3	0.21
Hazelnut	60.8	4.5	45.7	7.9	7.8	0.09
Peanut	49.2	6.8	24.4	15.8	15.6	0.00

Adopted from (Pradhan et al., nd) with slight modifications. All parameters are in g/100g. SFA means saturated fatty acids, MUFA means mono-unsaturated fatty acids and PUFA means poly-unsaturated fatty acids.

Phenolic compounds: Phytochemicals from tree nuts, including total phenols, flavonoids, proanthocyanidins (PAC), stilbenes, phytosterols, and carotenoids, exhibit diverse bioactivities such as antioxidant, antiviral, antiproliferative, hypocholesterolemic, and anti-inflammatory effects [47]. Polyphenolic compounds constitute the primary class of phytochemicals [48], with tree nuts serving as abundant sources of total phenolic compounds. Proanthocyanidins and hydrolysable tannins are generally the most abundantly found polyphenols in

some nuts (Table 2). Nuts also have a significant phytate content as well as flavonoid, phenolic acid and stilbene polyphenol classes [47]. It is widely recognized that the development of diabetic complications is influenced by various factors, with metabolic inflammation being a significant contributor to this process [13]. A growing body of evidence from clinical studies demonstrates the notable benefits of inflammation-targeted therapies in hyperglycemia, β -cell dysfunction, and insulin resistance in T2DM [13]. The pleiotropic benefits of phytochemicals

in diabetes include lowering LDL cholesterol as well as blood sugar, inhibiting inflammation, enhancing insulin

sensitivity, alleviating oxidative stress, offering vascular protection as well as reducing retinal inflammation [13].

Table 2. List of important phytochemicals found in nuts and their role in diabetes management.

Nuts	Phenolic compound	Potential health benefits in diabetes	References
Almond	Catechin, epicatechin, naringenin, eriodictyol, gallic acid, caffeic acid, chlorogenic acid, o-coumaric acid, p-coumaric acid, ferulic acid, hydroxybenzoic acid, protocatechuic, vanillic acid, quercetin, kaempferol, isorhamnetin	Antioxidant, Lipid lowering, hypoglycemic	[21,39,40,49]
Walnuts	Vanillic acid, catechin, pyrocatechin, protocatechuic acid, epicatechin, syringic acid, gallic acid, juglone and cinnamic acid, ellagic acid, rutin	Lipid lowering, anticholesteremic	[39,40,49]
Pistachio	Cyanidin, gallic acid, protocatechuic, eriodictyol, catechin, epicatechin, epicatechin gallate, luteolin, quercetin, myricetin, procyanidin B1, trimers, and tetramers	Lipid lowering, hypoglycaemic	[39,40,49]
Cashew	(+)- catechin, (-)-epicatechin, epigallocatechin and catechin gallate, syringic acid, gallic acid, delphinidin, cyanidin, quercetin, catechin, myricetin	Antiobesity, cardiovascular protection, antidiabetic	[39,40,50–53]
Pinenut	Catechin, epicatechin, vanillic acid, syringic acid, caffeic, hydroxycinnamic acid	Antioxidant, anti-inflammatory, antidiabetic, antihypertensive	[40,49]
Pecan	Ellagic acid, catechin, gallic acid, hydroxybenzoic acid, trans-cinnamic acid, syringic acid, caffeic acid, p-coumaric acid, ferulic acid, naringenin, apigenin, quercetin, rutin, kaempferol, isorhamnetin, resveratrol	Lipid lowering, hypoglycaemic	[39,40,49]
Macadamia	ampesterol, clerosterol, β -sitosterol, campestanol, Δ^5 -avenasterol + β -sitostanol, $\Delta^5,24(25)$ -stigmastadienol, Δ^7 -stigmastanol, 24- methylenecycloartenol, citrostadienol	Antioxidant, anti-dyslipidaemia, anti-inflammatory	[39,40,49,54]
Hazelnut	Gallic acid, protocatechuic acid, caffeic acid, o-coumaric acid, p-coumaric acid, ferulic acid, catechin, epicatechin, epicatechin gallate, rutin	Lipid lowering	[39,40,49]
Peanut	Catechin, epicatechin, quercetin, isorhamnetin, gallic acid, protocatechuic, caffeic acid, p-coumaric acid, procyanidins A and B, trimers and tetramers, prodelfinidin	Hypoglycaemic, antioxidant	[39,40,49]

Evidence of the impact of nuts on post prandial hyperglycaemia: Persistent elevation of blood sugar levels exerts detrimental effects on the initiation and advancement of both microvascular and macrovascular complications in individuals diagnosed with T2DM [55]. It has been demonstrated that postprandial hyperglycaemia (PPHG) may yield comparable or even more detrimental consequences than hyperglycaemia observed during fasting periods [56]. Previous

approaches to reduce PPHG in patients with diabetes have included improving insulin sensitivity (via exercise and weight loss), limiting total carbohydrate intake, and use of post-prandial glucose regulator drugs like Acarbose, Nateglinide, Repaglinide etc [55]. There is relatively poor long term adherence to most of these recommendations, additionally, drugs can be expensive for some patients, in some cases (20-30% of the users) the drugs have been shown to have adverse health

effects such as stomach aches, bloating and indigestion [55]. Tree nuts, on the other hand are a novel approach in the management of PPHG considering their nutrient and phytochemical composition [39,57].

The effects of nuts have been investigated in several studies. Gulati et al showed that dietary intervention with a premeal load (20g) of almonds (before breakfast, lunch and dinner daily) resulted in reduced postprandial glucose and insulin spikes, consequently mitigating the overall hyperglycaemic state in a cohort of Asian Indian individuals diagnosed with prediabetes[55]. Additionally, the intervention by Gulati et al led to a reversal to normal glucose regulation from the state of prediabetes. All these achievements were reached without restricting the carbohydrate intake of the participants meaning the almond nut can be used by diabetic patients to control blood glucose without the discomfort of altering their normal diet thus making it easier to incorporate as part of their lifestyle. Interestingly, the almond nuts used in Gulati et al's study did not have any adverse effects on the pre-diabetic participants, making it more superior and safer compared to the conventional postprandial glucose regulator drugs. The salutary effects of almond nuts on blood glucose level can be attributed to the MUFAs and soluble fiber in the nuts [57]. These have been shown to lead to the release of peptides such as GLP-1, GIP, and cholecystokinin (CCK) which can slow gastric emptying and stimulate insulin secretion [58]. Furthermore, the inclusion of almonds increases a feeling of satiety and leads to a strong dietary compensation effect[58]. In line with the above report, Li et al showed that almonds replacing 20% calories of the control diet improved glycemic control in Chinese T2DM patients, it however did not reduce glucose level to the unimpaired range (110-125 mg/dL [6.1- 6.9 mmol/L]) [59]. Another study by reported that the consumption of almonds with white bread was shown to significantly lower the postprandial area under the insulin concentration vs. time curve when compared to a high glycaemic index (GI)

meal (instant mashed potatoes) and significantly lowered the glucose peak height when compared with white bread, implying that, nuts consumption resulted in a lower postprandial hyperglycaemia[60]. Although almonds are rich in fat, they possess a low-glycemic index and could alter the glycemic index of co-consumed foods [59].

A comprehensive meta-analysis of twelve randomized controlled trials (RCTs) with a follow-up period exceeding three weeks indicates that the ingestion of an average daily quantity of 56 grams of tree nuts can enhance glycemic regulation among individuals diagnosed with Type 2 Diabetes (T2D) when compared to an isocaloric diet devoid of tree nuts [61]. Furthermore, findings from the investigation conducted by Kendall et al. underscore that the consumption of nuts either independently or in conjunction with a high glycemic index (GI) regimen contributes to the reduction of postprandial blood glucose levels[62]. Numerous dietary components found in nuts such as fiber and polyphenols, including both flavonoids and non-flavonoids, have been suggested to exhibit prebiotic properties and influence the regulation of glucose metabolism [63]. Studies have demonstrated that the alteration of gut microbiota through prebiotic interventions can enhance glycemic regulation in both healthy individuals and those with diabetes [64,65].

On the contrary, a study by Graveinstein et al, showed that long term consumption of almonds significantly decreased insulin sensitivity, and increased postprandial glucose concentrations in prediabetics [57]. The inconsistent reports regarding the hypoglycaemic effects of almonds can possibly be attributed to the different designs used in these studies. Additionally, almond dosage, background diet, and ethnicity might be accountable for the underlying factors for the discrepancy in the benefits of almond consumption in diabetes[59]. Nuts affect glycaemic response in a dose-dependent manner. The dose and duration of

supplementation in studies that observed a significant effect of nuts consumption on glycaemic factors ranged from 30–60 g/day and 4–24 weeks, respectively[66]. Although findings from a few earlier studies suggest a notable decrease in fasting insulin levels or enhancement in insulin resistance after the ingestion of nuts[66,67], later studies such as that of Palacios et al recently reported conflicting results regarding effect of nuts on insulin sensitivity [68] suggesting that there is still need for more studies to get more clarity on this matter.

Evidence of nut consumption on Metabolic Parameters:

It is widely agreed that cardiovascular diseases and diabetes share some risk factors such as increased adiposity and dyslipidaemia [69,70]. Thus, interventions aimed at preventing T2D should also target the cardiovascular-related risk factors such as dyslipidaemia. Several studies have reported the beneficial effects of nuts consumption on lipid profiles, especially total cholesterol (TC) and low-density lipoprotein cholesterol (LDL-C), in both healthy and high-cholesterolemic individuals from various geographical areas[66]. Nuts are rich in bioactive nutrients, such as γ -tocopherol, polyphenols, monounsaturated fatty acid (MUFA; oleic acid) and polyunsaturated fatty acid (PUFA; linoleic acid), known to have cardioprotective, antioxidant and lipid lowering properties[66]. Several studies have demonstrated these beneficial effects, for example, Cogan et al reported that pecan nuts significantly reduced post prandial blood lipids (total cholesterol, LDL cholesterol and non HDL cholesterol)[70]. Similarly, Jung et al showed that consumption of 56g of almonds for a period of 4 weeks decreased TC, LDL-C, and non-HDL-C among obese Korean adults compared to those on the control diet [71]. Several other randomised controlled trials reported the beneficial effects of consuming almonds on lipid profiles of either health or diabetic participants [72–76] even though they were done on different ethnic populations using different doses

(56g/day) and intervention periods (4–24weeks). Another study with pistachios revealed lipid lowering effects on TC as well as the ratio of total HDL-C and triglycerides on participants with well controlled T2DM [77]. Like almonds, pistachios have these lipid lowering effects in both healthy and diabetic individuals as shown by Gulati et al and Sari et al [78,79]. These observations imply that nuts can be used both in the prevention and management of diabetes by different ethnic groups. Other studies have shown that beyond lipid lowering, nuts have other favourable effects such as improving indices of inflammation and oxidative status [77,79]. Inflammation is a constant feature associated with the onset and progression of many chronic degenerative diseases, all three types of diabetes and one of the leading causes of its insurgence is the high adipose tissue content[80]. A study by Renzo et al showed that feeding 40g/day to healthy individuals over 4 weeks resulted in a significant up-regulation of sodium dismutase (SOD), catalase (CAT), peroxisome proliferator-activated receptor gamma (PPAR- γ), and Angiotensin I-converting enzyme (ACE) at the end of the study[81]. A clinical trial by Liu et al showed that almond diet (56 g/day) for 4 weeks reduced C-reactive protein (CRP) by a median 10.3 % (95% CI: –24.1, 40.5), tumor necrosis factor (TNF- α) by a median 15.7 % (95% CI: –0.3, 29.9), and interleukin-6 (IL-6) by a median 10.3 % (95% CI: 5.2, 12.6 %) in participants with T2DM in comparison to those in the control diet[82].

Mechanisms by which nuts ameliorate diabetes and metabolic syndrome:

Metabolic syndrome is characterized by central obesity, insulin resistance, dyslipidaemia, and hypertension, all of which are also key risk factors for Type 2 Diabetes Mellitus (T2DM) [7]. Nuts have shown unique potential in addressing multiple features of this syndrome through interconnected nutritional and biochemical pathways[16]. The diverse biochemical profiles of nuts confer unique health effects

through multifactorial mechanisms relevant to diabetes management [16,17]. For instance, walnuts are distinguished by their high α -linolenic acid (omega-3) content, which modulates inflammatory pathways via eicosanoid biosynthesis [16,17]. Almonds, rich in MUFAs and vitamin E, improve insulin sensitivity and exert antioxidative actions through the enhancement of glutathione peroxidase and catalase activity [17]. Pistachios, uniquely high in lutein and zeaxanthin, promote vascular health by upregulating endothelial nitric oxide synthase (eNOS), reducing oxidative stress and improving vasodilation [17]. Cashews, noted for their high magnesium and zinc levels, influence insulin receptor activity and β -cell preservation. Mechanistically, polyphenols in all nuts modulate AMP-activated protein kinase (AMPK) pathways, thus enhancing glucose uptake and fatty acid oxidation. Additionally, nut-derived dietary fiber modulates gut microbiota composition, leading to increased production of short-chain fatty acids (SCFAs) like butyrate, which has anti-inflammatory effects and improves insulin sensitivity [22].

- **Improvement of insulin sensitivity:** Monounsaturated and polyunsaturated fatty acids (especially oleic and linoleic acids in almonds, walnuts, and pistachios) upregulate PPAR- γ and downregulate inflammatory cytokines (e.g., TNF- α , IL-6), improving insulin signalling and β -cell function.
- **Modulation of postprandial glycaemia:** Soluble dietary fiber forms viscous gels that delay carbohydrate absorption, while phenolics and lipids in nuts enhance GLP-1 and CCK release, delaying gastric emptying and reducing glucose spikes.
- **Lipid-lowering and anti-atherogenic effects:** Phytosterols in nuts reduce intestinal cholesterol absorption, and their high

PUFA/MUFA content improves lipid profiles by lowering LDL and raising HDL cholesterol.

- **Reduction of inflammation and oxidative stress:** Polyphenols, flavonoids (like quercetin, catechin), and tocopherols act on NF- κ B and Nrf2 pathways to reduce systemic inflammation, a key driver of metabolic syndrome progression.
- **Gut microbiota modulation:** Prebiotic fibers and polyphenols from nuts alter gut microbial composition, increasing the production of short-chain fatty acids like butyrate, which improves insulin sensitivity and reduces systemic inflammation.

Together, these mechanisms explain why nuts are effective and how they intervene at multiple pathophysiological points relevant to both diabetes and metabolic syndrome. This multifaceted functionality makes nuts distinct among dietary interventions.

Safety considerations: Nut consumption can offer numerous health benefits due to their nutrient-rich profile, including healthy fats, protein, fiber, vitamins, and minerals. However, tree nuts have been reported to contain allergens such as seed storage proteins, pathogenesis-related (PR) proteins, profilins and lipid transfer proteins (LTP) [83] which may affect both children and adults. These allergens found in nuts can trigger allergic reactions ranging from mild to severe [84]. An allergic reaction to nuts can manifest as hives, itching, swelling, difficulty breathing, or even anaphylaxis, a life-threatening condition. Some individuals may experience adverse effects from consuming certain nuts due to natural toxins or anti-nutrients present in them. For instance, raw cashews contain urushiol, the same compound found in poison ivy, which can cause skin irritation or allergic reactions in susceptible individuals[85]. However, commercially available

cashews are typically heat-treated to remove this toxin. Food processing has been proposed as the method of choice to alter the allergenicity and toxicity of most foods including nuts to ensure their safety and improve their organoleptic properties [83,84]. While nuts offer numerous health benefits, excessive consumption can lead to nutritional imbalances. Nuts are energy-dense and high in fat, albeit predominantly healthy unsaturated fats. Overconsumption of nuts, especially if not balanced with other nutrient-rich foods, could contribute to weight gain or other health issues [86]. Lastly, whole nuts, particularly for young children or elderly individuals, can pose a choking hazard [87]. Chopping or grinding nuts before serving can help reduce the risk of choking

CONCLUSIONS

Nuts (mainly pistachios, almonds, and walnuts) have been shown to have beneficial effects on post prandial glycemic and lipid parameters. The higher monounsaturated fatty acid content and a lower polyunsaturated/saturated fat ratio reported in most edible nuts such as almond and pistachio nuts may be responsible for their favourable effect on lipids. Nuts can therefore be recommended as a dietary intervention for reversing the prediabetic stage and its related complications as well as reducing the progression to diabetes. While nuts offer numerous health benefits, individuals with nut allergies should take precautions to avoid exposure, and everyone should practice moderation and be mindful of portion sizes when incorporating nuts into their diet. There are fewer studies that have evaluated the effects of nuts on insulin resistance, and the few that are there report conflicting results. There is need for more studies to investigate the effects of nuts on insulin resistance in diabetic individuals. Although several studies have shown the pleiotropic benefits of nuts in diabetes, there is still a lack of clarity on the dose and duration of the intervention,

more research should focus on bring light on these issues to make it easier to incorporate nuts into as a dietary therapy.

Competing interest: The author has no financial interests or conflicts of interest.

Authors' contribution: The author wrote the entire review article.

Acknowledgments: There was no external funding supporting this publication.

REFERENCES

1. Almigbal TH, Alzarrah SA, Aljanoubi FA, Alhafez NA, Aldawsari MR, Alghadeer ZY, et al. Clinical Inertia in the Management of Type 2 Diabetes Mellitus: A Systematic Review. Vol. 59, Medicina (Lithuania). MDPI; 2023.
DOI: <https://doi.org/10.3390/medicina59010182>
2. Grasgruber P. Back to the pre-industrial age? FAOSTAT statistics of food supply reveal radical dietary changes accompanied by declining body height, rising obesity rates, and declining phenotypic IQ in affluent Western countries. Ann Med [Internet]. 2025 Dec 31;57(1). Available from: <https://www.tandfonline.com/doi/full/10.1080/07853890.2025.2514073>
3. Wu YL, Lin ZJ, Li CC, Lin X, Shan SK, Guo B, et al. Epigenetic regulation in metabolic diseases: mechanisms and advances in clinical study. Vol. 8, Signal Transduction and Targeted Therapy. Springer Nature; 2023.
DOI: <https://doi.org/10.1038/s41392-023-01333-7>
4. Siam NH, Snigdha NN, Tabasumma N, Parvin I. Diabetes Mellitus and Cardiovascular Disease: Exploring Epidemiology, Pathophysiology, and Treatment Strategies. Vol. 25, Reviews in Cardiovascular Medicine. IMR Press Limited; 2024.
DOI: <https://doi.org/10.31083/j.rcm2512436>.
5. Priyadarshini A, Madan R, Das S. Genetics and epigenetics of diabetes and its complications in India. Vol. 143, Human Genetics. Springer Science and Business Media Deutschland GmbH; 2024. p. 1–17.
DOI: <https://doi.org/10.1007/s00439-023-02616-3>
6. He KJ, Wang H, Xu J, Gong G, Liu X, Guan H. Global burden of type 2 diabetes mellitus from 1990 to 2021, with projections of prevalence to 2044: a systematic analysis across SDI levels

- for the global burden of disease study 2021. *Front Endocrinol (Lausanne)*. 2024;15.
DOI: <https://doi.org/10.3389/fendo.2024.1501690>
7. Jyotsna F, Ahmed A, Kumar K, Kaur P, Chaudhary MH, Kumar S, et al. Exploring the Complex Connection Between Diabetes and Cardiovascular Disease: Analyzing Approaches to Mitigate Cardiovascular Risk in Patients With Diabetes. *Cureus*. 2023 Aug 22.
DOI: <https://doi.org/10.7759/cureus.43882>
8. Avdic T, Carlsen HK, Rawshani A, Gudbjörnsdottir S, Mandalenakis Z, Eliasson B. Risk factors for and risk of all-cause and atherosclerotic cardiovascular disease mortality in people with type 2 diabetes and peripheral artery disease: an observational, register-based cohort study. *Cardiovasc Diabetol*. 2024 Dec 1;23(1).
DOI: <https://doi.org/10.1186/s12933-024-02226-x>
9. Zaninotto P, Steptoe A, Shim EJ. CVD incidence and mortality among people with diabetes and/or hypertension: Results from the English longitudinal study of ageing. *PLoS One*. 2024 May 1;19(5 May).
DOI: <https://doi.org/10.1371/journal.pone.0303306>
10. An Y, Xu B tuo, Wan S rong, Ma X mei, Long Y, Xu Y, et al. The role of oxidative stress in diabetes mellitus-induced vascular endothelial dysfunction. Vol. 22, *Cardiovascular Diabetology*. BioMed Central Ltd; 2023.
DOI: <https://doi.org/10.1186/s12933-023-01965-7>
11. Li Y, Liu Y, Liu S, Gao M, Wang W, Chen K, et al. Diabetic vascular diseases: molecular mechanisms and therapeutic strategies. Vol. 8, *Signal Transduction and Targeted Therapy*. Springer Nature; 2023.
DOI: <https://doi.org/10.1038/s41392-023-01400-z>
12. Jyotsna F, Ahmed A, Kumar K, Kaur P, Chaudhary MH, Kumar S, et al. Exploring the Complex Connection Between Diabetes and Cardiovascular Disease: Analyzing Approaches to Mitigate Cardiovascular Risk in Patients With Diabetes. *Cureus*. 2023 Aug 22.
DOI: <https://doi.org/10.7759/cureus.43882>
13. Kong M, Xie K, Lv M, Li J, Yao J, Yan K, et al. Anti-inflammatory phytochemicals for the treatment of diabetes and its complications: Lessons learned and future promise. Vol. 133, *Biomedicine and Pharmacotherapy*. Elsevier Masson s.r.l.; 2021. •
DOI: <https://doi.org/10.1016/j.biopha.2020.110975>
14. Nishi SK, Vigiouliou E, Kendall CWC, Jenkins DJA, Hu FB, Sievenpiper JL, et al. Nuts in the Prevention and Management of Type 2 Diabetes. Vol. 15, *Nutrients*. MDPI; 2023. DOI: <https://doi.org/10.3390/nu15040878>
15. Apostolopoulou M, Lambadiari V, Roden M, Dimitriadis GD. Insulin Resistance in Type 1 Diabetes: Pathophysiological, Clinical, and Therapeutic Relevance. Vol. 46, *Endocrine Reviews*. Endocrine Society; 2025. p. 317–48.
DOI: <https://doi.org/10.1210/endrev/bnae032>
16. Astbury NM. Interventions to improve glycaemic control in people living with, and at risk of developing type 2 diabetes. *Diabetes Obes Metab*. 2024 Sep 1;26(S4):39–49.
DOI: <https://doi.org/10.1111/dom.15855>
17. Arnesen EK, Thorisdottir B, Bärebring L, Söderlund F, Nwaru BI, Spielau U, et al. Nuts and seeds consumption and risk of cardiovascular disease, type 2 diabetes and their risk factors: a systematic review and meta-analysis. Vol. 67, *Food and Nutrition Research*. Swedish Nutrition Foundation; 2023.
DOI: <https://doi.org/10.29219/fnr.v67.8961>
18. Iizuka K, Yabe D. Dietary and Nutritional Guidelines for People with Diabetes. Vol. 15, *Nutrients*. Multidisciplinary Digital Publishing Institute (MDPI); 2023.
DOI: <https://doi.org/10.3390/nu15204314>
19. Mukwevho PL, Kaseke T, Fawole OA. Innovations in Biodegradable Packaging and Edible Coating of Shelled Temperate Nuts. *Food and Bioprocess Technology*. Springer; 2025. DOI: <https://doi.org/10.1007/s11947-025-03913-7>
20. Mirzaev Kamal Karimovich B, Uko P. THE POTENCY OF NUTS IN HUMAN HEALTH. Vol. 6, *ACADEMIC JOURNAL OF GLOBAL WHO IS WHO IN ACADEMIA*. NEW YORK CITY; 2025.
21. Gonçalves B, Pinto T, Aires A, Morais MC, Bacelar E, Anjos R, et al. Composition of Nuts and Their Potential Health Benefits—An Overview. Vol. 12, *Foods*. MDPI; 2023.
DOI: <https://doi.org/10.3390/foods12050942>
22. Rajaram S, Damasceno NRT, Braga RAM, Martinez R, Krishnatherton P, Sala-Vila A. Effect of Nuts on Markers of Inflammation and Oxidative Stress: A Narrative Review. Vol. 15, *Nutrients*. Multidisciplinary Digital Publishing Institute (MDPI); 2023. DOI: <https://doi.org/10.3390/nu15051099>
23. Mayomi PT, Aderinola TA. Proximate, Mineral, Amino Acid Composition, and Bioactive Properties of Dough Meals Supplemented with African Walnut Flour. *Food Science and Engineering [Internet]*. 2024 Oct 25;404–17.
DOI: <https://doi.org/10.37256/fse.5220245157>
24. Zhang S, Wang L, Fu Y, Jiang JC. Bioactive constituents, nutritional benefits and woody food applications of *Castanea mollissima*: A comprehensive review. Vol. 393, *Food Chemistry*. Elsevier Ltd; 2022.
DOI: <https://doi.org/10.1016/j.foodchem.2022.133380>

25. Siebeneichler TJ, Hoffmann JF, Galli V, Zambiasi RC. Composition and impact of pre- and post-harvest treatments/factors in pecan nuts quality. Vol. 131, Trends in Food Science and Technology. Elsevier Ltd; 2023. p. 46–60.
26. Santos MJ, Pinto T, Vilela A. Sweet Chestnut (*Castanea sativa* Mill.) Nutritional and Phenolic Composition Interactions with Chestnut Flavor Physiology. Vol. 11, Foods. MDPI; 2022. DOI: <https://doi.org/10.3390/foods11244052>
27. Liu J, Qin L, Zheng J, Tong L, Lu W, Lu C, et al. Research Progress on the Relationship between Vitamins and Diabetes: Systematic Review. Vol. 24, International Journal of Molecular Sciences. Multidisciplinary Digital Publishing Institute (MDPI); 2023. DOI: <https://doi.org/10.3390/ijms242216371>
28. Zhang T, O'Connor C, Sheridan H, Barlow JW. Vitamin K2 in Health and Disease: A Clinical Perspective. Vol. 13, Foods. Multidisciplinary Digital Publishing Institute (MDPI); 2024. DOI: <https://doi.org/10.3390/foods13111646>
29. Andr  s E, Lorenzo-Villalba N, Terrade JE, M  ndez-Bailon M. Fat-Soluble Vitamins A, D, E, and K: Review of the Literature and Points of Interest for the Clinician. Vol. 13, Journal of Clinical Medicine. Multidisciplinary Digital Publishing Institute (MDPI); 2024. DOI: <https://doi.org/10.3390/jcm13133641>
30.   wiel  g-Drabek M, Nie  -Le  niak J, Bialek-Dratwa A, Piekut A, Kiciak A, Dziubanek G, et al. Evaluation of Cadmium, Lead, Chromium, and Nickel Content in Various Types of Nuts: Almonds, Cashews, Hazelnuts, Peanuts, and Walnuts –Health Risk of Polish Consumers. Biol Trace Elem Res. 2024. DOI: <https://doi.org/10.1007/s12011-024-04438-4>
31. Liu Z, Li P, Zhao ZH, Zhang Y, Ma ZM, Wang SX. Vitamin B6 prevents endothelial dysfunction, insulin resistance, and hepatic lipid accumulation in Apoe ^{-/-} mice fed with high-fat diet. J Diabetes Res. 2016. DOI: <https://doi.org/10.1155/2016/1748065>
32. Mateos R, Salvador MD, Fregapane G, Goya L. Why Should Pistachio Be a Regular Food in Our Diet? Vol. 14, Nutrients. MDPI; 2022. • DOI: <https://doi.org/10.3390/nu14153207>
33. Mason SA, Keske MA, Wadley GD. Effects of Vitamin C Supplementation on Glycemic Control and Cardiovascular Risk Factors in People With Type 2 Diabetes: A GRADE-Assessed Systematic Review and Meta-analysis of Randomized Controlled Trials. Diabetes Care. 2021 Feb 1;44(2):618–30. DOI: <https://doi.org/10.2337/dc20-1893>
34. Tareke AA, Hadgu AA. The effect of vitamin C supplementation on lipid profile of type 2 diabetic patients: a systematic review and meta-analysis of clinical trials. Vol. 13, Diabetology and Metabolic Syndrome. BioMed Central Ltd; 2021. DOI: <https://doi.org/10.1186/s13098-021-00640-9>.
35. Tan SY, Dhillon J, Mattes RD. A review of the effects of nuts on appetite, food intake, metabolism, and body weight. In: American Journal of Clinical Nutrition. American Society for Nutrition; 2014. DOI: <https://doi.org/10.3945/ajcn.113.071456>
36. Markiewicz-  ukowska R, Pu  cion-Jakubik A, Grabia M, Perkowski J, Nowakowski P, Bielecka J, et al. Nuts as a Dietary Enrichment with Selected Minerals—Content Assessment Supported by Chemometric Analysis. Foods. 2022 Oct 1;11(20). DOI: <https://doi.org/10.3390/foods11203152>
37. Plessi M, Bertelli D, Monzani A, Simonetti - MS, Neri - A, Damiani - P. Dietary Fiber and Some Elements in Nuts and Wheat Brans [Internet]. Vol. 12, JOURNAL OF FOOD COMPOSITION AND ANALYSIS. 1999. DOI: <https://doi.org/10.1006/jfca.1999.0812>
38. Dhingra D, Michael M, Rajput H, Patil RT. Dietary fibre in foods: A review. Vol. 49, Journal of Food Science and Technology. 2012. p. 255–66. DOI: <https://doi.org/10.1007/s13197-011-0365-5>
39. Ros E. Health benefits of nut consumption. Vol. 2, Nutrients. MDPI AG; 2010. p. 652–82. DOI: <https://doi.org/10.3390/nu2070652>
40. Pradhan C, Peter N, Dileep N. Nuts as Dietary Source of Fatty Acids and Micro Nutrients in Human Health [Internet]. Available from: www.intechopen.com
41. Li X, Petrov MS. Dietary Fibre for the Prevention of Post-Pancreatitis Diabetes Mellitus: A Review of the Literature and Future Research Directions. Vol. 16, Nutrients. Multidisciplinary Digital Publishing Institute (MDPI); 2024. DOI: <https://doi.org/10.3390/nu16030435>
42. Abrams SA, Griffin IJ, Hawthorne KM. The Journal of Nutrition Inulin and Oligofructose: Health Benefits and Claims-A Critical Review Young Adolescents Who Respond to an Inulin-Type Fructan Substantially Increase Total Absorbed Calcium and Daily Calcium Accretion to the Skeleton 1-3. Vol. 137, J. Nutr. 2007. DOI: <https://doi.org/10.1093/jn/137.11.2524S>.
43. Tarfeen N, Nisa KU, Ahmad MB, Waza AA, Ganai BA. Metabolic and Genetic Association of Vitamin D with

- Calcium Signaling and Insulin Resistance. Vol. 38, Indian Journal of Clinical Biochemistry. Springer; 2023. p. 407–17. DOI: <https://doi.org/10.1007/s12291-022-01105-0>
44. Ko J, Kimita W, Skudder-Hill L, Li X, Priya S, Bharmal SH, et al. Dietary carbohydrate intake and insulin traits in individuals after acute pancreatitis: Effect modification by intra-pancreatic fat deposition. *Pancreatology*. 2021 Mar 1;21(2):353–62. DOI: <https://doi.org/10.1016/j.pan.2021.01.018>
45. Ağagündüz D, Icer MA, Yesildemir O, Koçak T, Kocigit E, Capasso R. The roles of dietary lipids and lipidomics in gut-brain axis in type 2 diabetes mellitus. Vol. 21, *Journal of Translational Medicine*. BioMed Central Ltd; 2023. DOI: <https://doi.org/10.1186/s12967-023-04088-5>
46. Bartelt A, Koehne T, Tödter K, Reimer R, Müller B, Behler-Janbeck F, et al. Quantification of bone fatty acid metabolism and its regulation by adipocyte lipoprotein lipase. *Int J Mol Sci*. 2017 Jun 13;18(6). DOI: <https://doi.org/10.3390/ijms18061264>
47. Alasalvar C, Bolling BW. Review of nut phytochemicals, fat-soluble bioactives, antioxidant components and health effects. Vol. 113, *British Journal of Nutrition*. Cambridge University Press; 2015. p. 568–78. DOI: <https://doi.org/10.1017/S0007114514003729>
48. Martirosyan D. Functional Food Science and Bioactive Compounds. *BCHD Bioactive Compounds in Health and Disease* [Internet]. 2025;8(6):218–29. DOI: <https://www.doi.org/10.31989/bchd.v4i3.786>
49. Woźniak M, Waśkiewicz A, Ratajczak I. The Content of Phenolic Compounds and Mineral Elements in Edible Nuts. *Molecules*. 2022 Jul 1;27(14). DOI: <https://doi.org/10.3390/molecules27144326>
50. Sruthi P, Naidu MM. Cashew nut (*Anacardium occidentale* L.) testa as a potential source of bioactive compounds: A review on its functional properties and valorization. Vol. 3, *Food Chemistry Advances*. Elsevier Ltd; 2023. DOI: <https://doi.org/10.1016/j.focha.2023.100390>
51. Chandrasekara N, Shahidi F. Antioxidative potential of cashew phenolics in food and biological model systems as affected by roasting. *Food Chem*. 2011 Dec 15;129(4):1388–96. DOI: <http://dx.doi.org/10.1016/j.foodchem.2011.05.075>
52. Sruthi P, Roopavathi C, Madhava Naidu M. Profiling of phenolics in cashew nut (*Anacardium occidentale* L.) testa and evaluation of their antioxidant and antimicrobial properties. *Food Biosci*. 2023 Feb 1;51. DOI: <https://doi.org/10.1016/j.fbio.2022.102246>
53. Chen YY, Li N yang, Guo X, Huang H jie, Garcia-Oliveira P, Sun J yue, et al. The nutritional and bio-active constituents, functional activities, and industrial applications of cashew (*Anacardium occidentale*): A review. *Food Front*. 2023 Dec 1;4(4):1606–21. <https://iadns.onlinelibrary.wiley.com/doi/epdf/10.1002/fft.2.250>
54. Insanu M, Hartati R, Bajri F, Fidrianny I. *Macadamia* genus: An updated review of phytochemical compounds and pharmacological activities. Vol. 11, *Biointerface Research in Applied Chemistry*. AMG Transcend Association; 2021. p. 14480–9. DOI: <https://doi.org/10.33263/BRIAC116.1448014489>
55. Gulati S, Misra A, Tiwari R, Sharma M, Pandey RM, Upadhyay AD, et al. Premeal almond load decreases postprandial glycaemia, adiposity and reversed prediabetes to normoglycemia: A randomized controlled trial. *Clin Nutr ESPEN*. 2023 Apr 1;54:12–22. DOI: <https://doi.org/10.1016/j.clnesp.2022.12.028>
56. Moon J, Kim JY, Yoo S, Koh G. Fasting and postprandial hyperglycemia: Their predictors and contributions to overall hyperglycemia in Korean patients with type 2 diabetes. *Endocrinology and Metabolism*. 2020;35(2):290–7. DOI: <https://doi.org/10.3803/EnM.2020.35.2.290>
57. Gravestijn E, Mensink RP, Plat J. The effects of long-term almond consumption on whole-body insulin sensitivity, postprandial glucose responses, and 48 h continuous glucose concentrations in males and females with prediabetes: a randomized controlled trial. *Eur J Nutr*. 2023 Sep 1;62(6):2661–72. DOI: <https://doi.org/10.1007/s00394-023-03178-w>
58. Kien CL. *Dietary Interventions for Metabolic Syndrome: Role of Modifying Dietary Fats*. 2009. DOI: <https://doi.org/10.1007/s11892-009-0009-6>
59. Li SC, Liu YH, Liu JF, Chang WH, Chen CM, Chen CYO. Almond consumption improved glycemic control and lipid profiles in patients with type 2 diabetes mellitus. *Metabolism*. 2011 Apr;60(4):474–9. DOI: <https://doi.org/10.1016/j.metabol.2010.04.009>
60. Jenkins DJA, Kendall CWC, Josse AR, Salvatore S, Brighenti F, Augustin LSA, et al. The Journal of Nutrition Nutrient Physiology, Metabolism, and Nutrient-Nutrient Interactions Almonds Decrease Postprandial Glycemia, Insulinemia, and

- Oxidative Damage in Healthy Individuals 1. Vol. 136, J. Nutr. 2006. DOI: <https://doi.org/10.1093/jn/136.12.2987>
61. Viguioliouk E, Kendall CWC, Mejia SB, Cozma AI, Ha V, Mirrahimi A, et al. Effect of tree nuts on glycemic control in diabetes: A systematic review and meta-analysis of randomized controlled dietary trials. Vol. 9, PLoS ONE. Public Library of Science; 2014. DOI: <https://doi.org/10.1371/journal.pone.0103376>
62. Kendall CWC, Esfahani A, Josse AR, Augustin LSA, Vidgen E, Jenkins DJA. The glycemic effect of nut-enriched meals in healthy and diabetic subjects. Nutrition, Metabolism and Cardiovascular Diseases. 2011 Jun;21(SUPPL. 1). DOI: <https://doi.org/10.1016/j.numecd.2011.03.013>
63. Kim YA, Keogh JB, Clifton PM. Polyphenols and glycémie control. Vol. 8, Nutrients. MDPI; 2016. DOI: <https://doi.org/10.3390/nu8010017>
64. Sadagopan A, Mahmoud A, Begg M, Tarhuni M, Fotso M, Gonzalez NA, et al. Understanding the Role of the Gut Microbiome in Diabetes and Therapeutics Targeting Leaky Gut: A Systematic Review. Cureus. 2023. DOI: <https://doi.org/10.7759/cureus.41559>
65. Crudele L, Gadaleta RM, Cariello M, Moschetta A. Gut microbiota in the pathogenesis and therapeutic approaches of diabetes. Vol. 97, eBioMedicine. Elsevier B.V.; 2023. DOI: <https://doi.org/10.1016/j.ebiom.2023.104821>
66. Khalili L, A-Elgadir TME, Mallick AK, El Enshasy HA, Sayyed RZ. Nuts as a Part of Dietary Strategy to Improve Metabolic Biomarkers: A Narrative Review. Vol. 9, Frontiers in Nutrition. Frontiers Media S.A.; 2022. DOI: <https://doi.org/10.3389/fnut.2022.881843>
67. Hernández-Alonso P, Salas-Salvadó J, Baldrich-Mora M, Juanola-Falgarona M, Bulló M. Beneficial effect of pistachio consumption on glucose metabolism, insulin resistance, inflammation, and related metabolic risk markers: A randomized clinical trial. Diabetes Care. 2014 Nov 1;37(11):3098–105. DOI: <https://doi.org/10.2337/dc14-1431>
68. Palacios OM, Maki KC, Xiao D, Wilcox ML, Dicklin MR, Kramer M, et al. Effects of Consuming Almonds on Insulin Sensitivity and Other Cardiometabolic Health Markers in Adults With Prediabetes. J Am Coll Nutr. 2020 Jul 3;39(5):397–406. DOI: <https://doi.org/10.1080/07315724.2019.1660929>
69. Forbes JM, Cooper ME. Mechanisms of Diabetic Complications. Physiol Rev [Internet]. 2013;93:137–88. DOI: <https://doi.org/10.1152/physrev.00045.2011>
70. Cogan B, Pearson RC, Paton CM, Jenkins NT, Cooper JA. Pecan-enriched diet improves cholesterol profiles and enhances postprandial microvascular reactivity in older adults. Nutrition Research. 2023 Mar 1;111:44–58. DOI: <https://doi.org/10.1016/j.nutres.2023.01.001>
71. Jung H, Chen CYO, Blumberg JB, Kwak HK. The effect of almonds on vitamin E status and cardiovascular risk factors in Korean adults: a randomized clinical trial. Eur J Nutr. 2018 Sep 1;57(6):2069–79. DOI: <https://doi.org/10.1007/s00394-017-1480-5>
72. Foster GD, Shantz KL, Vander Veur SS, Oliver TL, Lent MR, Virus A, et al. A randomized trial of the effects of an almond-enriched, hypocaloric diet in the treatment of obesity. American Journal of Clinical Nutrition. 2012 Aug 1;96(2):249–54. DOI: <https://doi.org/10.3945/ajcn.112.037895>
73. Liu Y, Hwang HJ, Ryu H, Lee YS, Kim HS, Park H. The effects of daily intake timing of almond on the body composition and blood lipid profile of healthy adults. Nutr Res Pract. 2017 Dec 1;11(6):479–86. DOI: <https://doi.org/10.4162/nrp.2017.11.6.479>
74. Berryman CE, West SG, Fleming JA, Bordi PL, Kris-Etherton PM. Effects of daily almond consumption on cardiometabolic risk and abdominal adiposity in healthy adults with elevated LDL-cholesterol: A randomized controlled trial. J Am Heart Assoc. 2015;4(1). DOI: <https://doi.org/10.1161/jaha.114.000993>
75. Sabaté J, Haddad E, Tanzman JS, Jambazian P, Rajaram S. Printed in USA. Vol. 77, Am J Clin Nutr. 2003.
76. Liu Y, Hwang HJ, Kim HS, Park H. Time and Intervention Effects of Daily Almond Intake on the Changes of Lipid Profile and Body Composition among Free-Living Healthy Adults. J Med Food. 2018 Apr 1;21(4):340–7. DOI: <https://doi.org/10.1089/jmf.2017.3976>
77. Sauder KA, McCrea CE, Ulbrecht JS, Kris-Etherton PM, West SG. Effects of pistachios on the lipid/lipoprotein profile, glycemic control, inflammation, and endothelial function in type 2 diabetes: A randomized trial. Metabolism. 2015 Nov 1;64(11):1521–9. DOI: <https://doi.org/10.1016/j.metabol.2015.07.021>
78. Gulati S, Misra A, Pandey RM, Bhatt SP, Saluja S. Effects of pistachio nuts on body composition, metabolic, inflammatory and oxidative stress parameters in Asian

- Indians with metabolic syndrome: A 24-wk, randomized control trial. *Nutrition*. 2014 Feb;30(2):192–7.
DOI: <https://doi.org/10.1016/j.nut.2013.08.005>
79. Sari I, Baltaci Y, Bagci C, Davutoglu V, Erel O, Celik H, et al. Effect of pistachio diet on lipid parameters, endothelial function, inflammation, and oxidative status: A prospective study. *Nutrition*. 2010 Apr;26(4):399–404.
DOI: <https://doi.org/10.1016/j.nut.2009.05.023>
 80. Calder PC, Ahluwalia N, Brouns F, Buetler T, Clement K, Cunningham K, et al. Dietary factors and low-grade inflammation in relation to overweight and obesity. *British Journal of Nutrition*. 2011;106(SUPPL. 3).
DOI: <https://doi.org/10.1017/S0007114511005460>
 81. Renzo L Di, Cioccoloni G, Bernardini S, Abenavoli L, Aiello V, Marchetti M, et al. A hazelnut-enriched diet modulates oxidative stress and inflammation gene expression without weight gain. *Oxid Med Cell Longev*. 2019;2019.
DOI: <https://doi.org/10.1155/2019/4683723>.
 82. Liu JF, Liu YH, Chen CM, Chang WH, Chen CYO. The effect of almonds on inflammation and oxidative stress in Chinese patients with type 2 diabetes mellitus: A randomized crossover controlled feeding trial. *Eur J Nutr*. 2013 Apr;52(3):927–35.
DOI: <https://doi.org/10.1007/s00394-012-0400-y>
 83. Cuadrado C, Sanchiz Á, Linacero R. *Nut Allergenicity: Effect of Food Processing*. 2021.
DOI: <https://doi.org/10.3390/allergies1030014>
 84. Astuti RM, Palupi NS, Suhartono MT, Kusumaningtyas E, Lioe HN. Effect of processing treatments on the allergenicity of nuts and legumes: A meta-analysis. *J Food Sci*. 2023 Jan 1;88(1):28–56.
DOI: <https://doi.org/10.1111/1750-3841.16381>
 85. An N, Pourzal S, Luccioli S, Vukmanović S. Effects of diet on skin sensitization by nickel, poison ivy, and sesquiterpene lactones. Vol. 137, *Food and Chemical Toxicology*. Elsevier Ltd; 2020.
 86. Godswill C, Somtochukwu V. NUTRITIONAL DISEASES AND NUTRIENT TOXICITIES: A SYSTEMATIC REVIEW OF THE DIETS AND NUTRITION FOR PREVENTION AND TREATMENT. Vol. ISSN, *International Journal of Advanced Academic Research | Sciences*. 2020.
 87. Ranjous Y, Al Balkhi A, Alnader I, Rkab M, Ataya J, Abouharb R. Knowledge and misconceptions of choking an[4]. The rise in the number of people with type 2 diabetes is of socio-economic, demographic, envirod first-aid procedures

among Syrian adults: A cross-sectional study. *SAGE Open Med*. 2024 Jan 1;12.